

Supporting farmers in the transition to agroecology to promote carbon sequestration from silvopastoral systems

Sara Burbi^a, Rodrigo Olave^b

^aCentre for Agroecology, Water and Resilience (CAWR), Coventry University, Ryton Organic Gardens, Wolston Lane, Ryton-on-Dunsmore CV8 3LG, England, United Kingdom, sara.burbi@coventry.ac.uk

^bAgri-Food and Biosciences Institute (AFBI), Sustainable Agri-Food Sciences Division, Large Park, Hillsborough BT26 6DR, Northern Ireland, United Kingdom, rodrigo.olave@afbini.gov.uk

Abstract: *Silvopastoral systems, or silvopastures, have the potential to provide a wide range of ecosystem services to benefit the environment and society in relation to climate change, in particular carbon sequestration. However, being considered highly diversified and integrated systems, silvopastures present several challenges in the evaluation of their performance to support their adoption in different environmental contexts with the purpose of increasing carbon sequestration from agroforestry. While such evaluations and predictions of carbon sequestration potential can be achieved by researchers with long-term experimental studies and scenario-based modelling, farmers and land managers may need sufficiently accurate yet simplified methodologies to estimate the impact of land use and land use changes from forestry (LULUCF). This can be achieved with decision-support tools and participatory work where farmers and researchers engage in sharing and co-creating knowledge on best agroecological practices. This work sets out to review the current knowledge on carbon sequestration from silvopastures and the state of the agroforestry sector in the UK. It then proposes a methodology to integrate agroecological practices and knowledge generated using scenario-based LULUCF modelling to identify practical farm management strategies. It highlights potential barriers to and drivers for innovation in the UK agroforestry sector, including the impact of farmers' attitudes to climate change and silvopasture, and the impacts of networks of influence, community of practice, and the current provision of extension and advisory services. Finally, it illustrates how a decision-support system tailored for the agroforestry sector can benefit farmers in their transition to silvopastoral farming systems by adopting agroecological principles.*

Keywords: *silvopasture, carbon sequestration, land use changes, farmer engagement, decision support tools, knowledge integration, agroecological transition*

Introduction

In the United Kingdom (UK), agriculture accounts for about 7% of the total greenhouse gas (GHG) emissions (DEFRA, 2016) and the country is committed to reduce these emissions by 80% below the 1990 value (United Kingdom Parliament, 2008). The removal of GHG emissions, especially CO₂ from the atmosphere due to land-use practices could potentially be used as a mitigation option by implementing agroecological systems. Agroecology is defined as the discipline that addresses practical aspects of resilient food production and natural resources management, their environmental impact as well as the governance and socio-economic challenges facing current food and farming systems (Altieri, 2002; Wezel et al., 2009; TWN and SOCLA, 2015). High-performing agroecological systems have the potential to ensure productivity and biodiversity by adopting climate-friendly practices (Pretty et al., 2011; Bohan et al., 2013). Agroecological practices favour the protection of soil carbon and carbon sequestration in vegetation biomass and therefore have great potential for climate change mitigation (Wibbelmann et al., 2013), significantly reducing the carbon footprint especially in smallholder farming (Rakotovo et al., 2017). Agroforestry can be considered among agroecological systems because it uses complementarities and synergies combining crops, plants, trees and animals within diverse spatial and temporal settings

(Altieri et al., 2015). Silvopasture, which is a type of agroforestry system that combines grazing stock and trees on the same land base, can provide high biodiversity, efficient feed conversion and good animal welfare, and mitigate soil erosion and climate change (Broom et al., 2013). However, when considering scale, the level of system change that would be needed to implement agroecological practices in agroforestry systems is high and the uptake of such systems is currently low in Europe (Wezel et al., 2014). This is also due to the complexity and long-term risk management needed in such highly diversified farming systems and difficulties in integrating scientific advances and practical support for farmers to design and implement silvopastoral practices (Lovell et al., 2010; Torralba, et al., 2016).

The three dimensions of agroecology, i.e. movement, science, practice, described by Wezel et al. (2009) are well reflected in the recent interest in silvopastoral systems and, more widely, in agroforestry to promote diversified and sustainable farming that contributes to mitigate climate change. Carbon sequestration from agriculture and agroforestry further contributes to reducing the carbon footprint of these production systems. However, challenges remain in the improvement of the accuracy of carbon accounting in complex systems, and the subsequent impact on promoting and supporting the transition to agroecological practices, as well as integrating these into agroforestry-adapted future policy. In order to understand how scientists can support farmers and land managers in the transition to agroecology to promote carbon sequestration in silvopastoral systems, we need to acknowledge the contribution of agroforestry, the methodologies used to estimate the impact from Land Use and Land Use Changes from Forestry (LULUCF), and the potential of agroecological practices to increase carbon sequestration from silvopasture.

Carbon sequestration from agroforestry (UK)

The role of land use systems in stabilizing the CO₂ levels and increasing the carbon (C) sink potential has attracted considerable scientific attention, especially after the Kyoto Protocol to the United Nations Framework Convention on Climate Change (UNFCCC, 2015). The UK reports GHG emissions and carbon stock changes for the LULUCF sector as part of its national GHG inventory. The LULUCF sector is unusual in that it is the only sector of the inventory which can report removals as well as emissions for GHGs. LULUCF removals occur when the carbon stocks of soils and biomass increase, while emissions occur when carbon stocks decrease, or when the non-carbon dioxide GHGs methane and nitrous oxide are released from soils or biomass as a result of change in land use or management or fires (DEFRA, 2016).

Agroforestry is one of the most important land use systems practised in diverse ecoregions around the world and have a special relevance within agroecology. These woody perennial-based land use systems have relatively high capacities for capturing and storing atmospheric CO₂ in vegetation, soils, and biomass products. Agroforestry in the UK is developing along two lines: silvoarable systems where rows of trees are intercropped with arable crops and silvopastoral systems where stock graze on pasture between widely spaced trees. Although both systems are not yet practised in a great extent, its role in reducing C emissions and promoting carbon sequestration has been recognised through various policy schemes at national and European level (Fornara et al., 2017; Hart and Baldock, 2011).

Silvopastoral systems have the potential to provide a wide range of ecosystem services to benefit the environment and society in relation to climate change, in particular carbon sequestration (Cardinael et al., 2017; Kumar and Nair, 2011; Olave, 2016; Wilson & Lowell, 2016). In order to promote the adoption of farming practices that increase carbon sequestration at farm level, we need to provide farmers and land managers with tools that are user-friendly and scientifically accurate. This work sets out to highlight possible obstacles in the transition to agroecology in silvopastoral systems and proposes a methodology to support farmers in their transition by adopting a practical focus grounded on improved carbon accounting modelling.

Potential barriers to transition

National Adaptation Strategies to address the impacts of climate change are implemented in all European members states, but results may vary, in particular in highly diversified sectors such as the agriculture and forestry. Challenges include the uncertainty regarding scientific knowledge on carbon balance and accounting methodologies, coupled with the difficulty in achieving successful multi-actor approach activities involving government agencies, local agencies, researchers and private sector, and in establishing an effective knowledge transfer network (Biesbroek et al., 2010). Long et al. (2016, p.17) identified a series of barriers to the adoption of climate-smart agricultural (CSA) technology. These include: *“low awareness of CSA and inaccessible language, high costs and long return on investment (ROI) periods, lack of verified impact of technologies, regulatory and policy issues, hard to reach and train farmers, Research and Development (R&D) and policies do not match to 'on-the-ground' reality, low consumer demand, and unequal distribution of costs/benefits across supply chains.”*

Furthermore, supporting farmers in the transition to silvopastoral systems can face obstacles linked to the potential lack of trust in government advice and reluctance in taking a financial risk, and the fragmentation of extension services provided in the United Kingdom. Lacking in trust of scientific evidence regarding climate change and carbon accounting can be reflected on difficulties for farmers and land managers to access scientific knowledge and body of academic literature on climate studies, largely inaccessible to the general public, and the perceived lack of transparency and clarity in such information, e.g. adopting overly technical terminology (Hofmann et al., 2011). Although there have been significant steps in promoting access to scientific knowledge to the wider public, e.g. Open Access publications, knowledge brokerage events, transition in farming systems can be seen as a great financial risk and motivations to adopt innovative practices in the farming sector can be primarily economic, then followed by the improvement in management practices, and then market pressure (Barnes et al., 2010). In the context of whether the financial burden is real or perceived, practitioners need tools that are specifically adapted to agroforestry which easily identify practices and economic trade-offs to support them in the decision-making process. However, it is difficult to include socio-economic trade-offs because they can be very context-specific in the farming sector and in particular in agroforestry, which is a highly diversified type of farming (Paracchini et al., 2008; de Boer et al., 2011; Vellinga et al., 2011, Wallach et al., 2016). The majority of studies on impact assessment and indicators used in agroforestry do not include economic variables (Fargerholm et al., 2016).

Promoting the adoption of agroecological practices in silvopastoral systems can have challenges associated with the promotion of innovation more generally. Farmers' attitudes to innovation can be influenced by several factors, such as personal beliefs, family values, and other external actors like extension officers, experts and also the media. In a recent British study on livestock farmers' attitudes to climate change, Burbi et al. (2016, p.467) found that a *“proactive attitude seems to be hindered by confusion and lack of confidence in governmental strategies to disseminate scientific knowledge.”* While extension officers provide specific, technical advice to farmers and land managers (Takemura et al., 2014), the funding and support to advisory and extension services in the United Kingdom have been progressively and continuously reduced in the past 3 decades, creating a highly fragmented sector with great variability in the efficiency and impact of these services (Swanson and Rajalahti, 2010). Studies suggest that focusing exclusively on technical advice may not be effective in promoting innovation and transition in farming (Islam et al., 2013; Llewellyn, 2007). Advisory, extension services and knowledge transfer activities need to include information on cost-effectiveness and estimates of the socio-economic impact of changes in farming practices (Kings and Ilbery, 2010) and need to take into account farmers' attitudes and perceptions of innovation (Mills et al., 2013). This is particularly relevant when addressing carbon sequestration and climate change related topics because of the difference in aims and objectives between researchers, farmers and land managers, and policy-makers, resulting in potential uncertainty over the long-term impact of the transition and the best risk management option farmers ought to take (Burbi and Olave, 2017).

Proposed methodology

Among the advantages of agroecology are its adaptability, site-specificity and the fact that it promotes the integration of scientific and traditional knowledge (Sarandón and Flores, 2014). These three characteristics make agroecology likely to be taken up by farmers (Saj et al., 2017). Participatory Action Research (PAR) is widely accepted by the scientific community as a multi-actor approach to engage with different groups of stakeholders with differing aims and knowledge(s) (IFAD, 2009; Pretty and Buck, 2002). This approach is particularly effective in agriculture and forestry extension services because it fosters improved communication and knowledge integration, leading to a collective process with practical problem-solving activities (German et al., 2012; Klerkx et al. 2012; Le Gal et al., 2011; Xu et al., 2011; Jacobi et al., 2016). In spite of the possible biases and limitations listed by Neef and Neubert (2011), e.g. context-dependent results, differing goals, power relationships, PAR creates a process that allows for greater engagement with stakeholders (Mapfumo et al., 2013; Oliver et al., 2012). In the case of farmers, the process helps to build trust and they are then more likely to adopt innovative practices as a result from improved communication with researchers and other experts, in particular when considering the economic impact of transition and the potential barriers represented by the lack of an agroforestry-adapted environmental policy and long-term political strategy integrating agriculture, agroforestry and climate change mitigation (den Herder et al., 2017; Emery and Franks, 2012; Mosquera-Losada et al., 2015).

The methodology we propose is grounded in the concept of knowledge integration as described by Newell et al. (2005) and the provision of evidence-based scientific advice using farmer-friendly indicators to design a decision support tool that provides practical advice to farmers and land managers wanting to improve carbon sequestration from silvopasture or transition from grassland-based to agroforestry systems. The framework is structured in three phases: i) Initially, more comprehensive and accurate LULUCF models can provide the evidence-base to support the transition to agroforestry systems with greater carbon sequestration potential; ii) Subsequently, specific silvopastoral practices that increase carbon sequestration can be selected based on the following criteria: agroecological principles, potential to sequester carbon and confidence of evidence-base to support it; iii) Finally, user-friendly indicators can be identified through participatory work with farmers and land managers to design and implement a decision support tool adapted to silvopasture and carbon sequestration. This last phase will lead to the possibility of expanding the framework to integrate a science-based farm management assessment and monitoring with economic and legislative contexts.

Phase 1: Improvement of LULUCF accounting could provide more accurate estimates of the potential for carbon sequestration based on land types and land use. On the one hand, this would allow for more accurate prediction of future scenarios in case of land use changes, providing evidence to support policy making (Rittenhouse and Rissman, 2012). On the other hand, this would also allow for the design and implementation of decision-support tools that are more accurate and better reflect the impact of agroecological land management practices on carbon sequestration (Mosquera-Losada et al., 2017). Recent studies have shown the benefit to GHG mitigation of combining tree, hedgerows and farming production in specific systems using life cycle assessment techniques (Nguyen et al., 2013, Black et al., 2014). However, as an example, some activities not included in the GHG inventory and LULUCF accounting are fences replaced by hedges, development of scrub, hedge management and C dynamics in soils below hedges. Therefore, this phase is expected to generate data on future scenarios based on more accurate LULUCF accounting of carbon balances under different land managements. This work is essential to bring clarity using appropriate and standardised agroforestry descriptions (Palma et al., 2015) and including economic considerations into the definition of agroforestry categories with the purpose to build more accurate scenarios relevant to the sector and, therefore, more likely to be useful from a practical, farmer's perspective (Keesman et al., 2011; Luedeling et al., 2014).

Phase 2: The five main agroecological principles for sustainable livestock farming are: 1) integrated animal health management; 2) reducing external inputs; 3) re-coupling carbon (C), nitrogen (N) and phosphorous (P) cycles; 4) preserving and using biodiversity; and 5) increasing systems diversity and resilience (Dumont et al., 2014). This phase will list specific agroecological practices in silvopasture and assign scores based on their potential to sequester carbon and on-farm empirical studies. As examples, soil carbon stocks at 1 m depth tend to be greater under the tree canopy, with 50.2 Mg ha⁻¹ at 2 m from cork oak and 26.5 Mg ha⁻¹ at 15 m (Howlett et al., 2011). Soil aggregates also play a significant role in C stocks in silvopastoral systems, where a greater proportion of micro-aggregates are found. Micro-aggregates have been found to be more stable than macro-aggregates in retaining carbon. Therefore, it is reasonable to expect that carbon sequestered in micro-aggregates in soil managed under silvopastoral farming conditions is more stable and likely to contribute to long-term C stocks. However, soil composition and C saturation rates also need to be considered (Fornara et al., 2017). A review of all agroecological practices in silvopasture that increase carbon sequestration will allow the initial assessment of the system based on criteria, such as soil characteristics, tree density, stocking rate, grazing patterns and pasture composition.

Phase 3: An indicator-based decision support tool will be designed based on data obtained from Phases 1 and 2 to provide farmers and land managers with practical solutions to implement and monitor their transition to more carbon-friendly systems. Sustainability assessment tools can adopt a variety of methodologies, e.g. indicators and indices, product-related assessment, integrated assessment (Ness et al., 2007). The challenges in this case are related to time and resources available to assess and monitor the farm, and the diversity of systems in terms of production, location, soil properties, and climatic variations. Therefore, rapid indicator-based assessment tools are the preferred methodology to allow for a context-specific assessment that is both accurate and user-friendly (Burbi et al., 2016). Indicator-based tools are supported by scientific evidence, yet easy-to-use, e.g. soil health indicators (Nicholls et al., 2014). They allow for regular and context-specific regular assessment of the system (Mwongera et al., 2017) and as a result they have the potential to promote greater researcher-farmer engagement and aid long-term evaluation and monitoring of the carbon balance in silvopastures. This approach is in line with what suggested by Csikvari et al. (2017) in a recent review of tools available for the design and management of agroforestry systems, which highlighted the lack of knowledge by farmers of the wider benefits of agroforestry, including carbon sequestration. The authors also highlight the difficulty in matching scientific knowledge with current ecological and economic needs of farmers and land managers and how tools need to support farmers in the identification of key practices to implement on-farm, being user-friendly and aimed at the adoption of a more strategic long-term approach to farm management to increase carbon sequestration from the whole system.

Advantages

Drivers for transition

Knowledge integration is key to the promotion of innovation among farmers. However, this process can be difficult to achieve because farmers' knowledge tends to be based on anecdotal evidence and practical experience and therefore it is difficult to integrate with scientific knowledge (Raymond et al., 2010; Sutherland et al., 2013). Participatory learning processes where farmers' knowledge is valued have the potential to foster on-farm innovation and increase the adoption of agroecological practices (Louah et al., 2017). The methodology described aims at fostering knowledge transfer between researchers and farmers, effectively creating synergies between the different actors involved. On the one hand, scientific knowledge can be communicated effectively to farmers and land managers who, on the other hand, will have the opportunity to contribute by feeding back on the most successful implementation strategies to promote carbon sequestration by the transition to silvopasture, valuing farmers' knowledge and contribution to the process (Schöll and Binder, 2009; Virji et al., 2012; Castellanos et al., 2013). Farmers tend to rely on peer-to-peer

exchange of experiences to acquire knowledge and therefore, the networks of influence in which farmers operate have a great impact on the adoption of innovation (Oreszczyn et al., 2010; McKenzie, 2011; Schut et al, 2015). By promoting collaboration between researchers and farmers, the methodology proposed is expected to be more successful in transferring knowledge to support the transition to a more sustainable system, without assuming that financial incentives will be the only effective drivers for change (Sereke et al., 2016). It represents a means to foster farmer-driven innovation capitalising on the strengthening of synergies with extension services and researchers.

Perceptions vs Reality

In spite of the growing interest in carbon sequestration from the farming and agroforestry sectors, farmers' attitudes to climate change related issues could be influenced greatly by their perceptions of risk and of clear long-term benefits from the transition to silvopasture. Farmers appreciate flexibility in the management of their agri-businesses and the policies that impact the sector (Jones et al., 2013). However, they are also more likely to adopt innovative practices that have a perceived low economic risk or have multiple benefits (Burbi et al., 2016; Khatri-Chhetri et al., 2017), especially when clear short-term benefits and income generation are presented, rather than long-term impacts (Tittonell et al., 2012). This may create a challenge in the design of decision-support tools that are accurate, yet flexible and comprehensive, address carbon sequestration from the environmental, socio-economic, and potentially also legislative point of view, as shown by studies on complex socio-ecological systems (Bodin and Tengö, 2010; Feola and Binder, 2010; Cornell et al., 2013). The proposed methodology focuses on evidence-base practical advice on silvopastoral systems using a farmer-friendly tool to identify the practices more beneficial in terms of carbon sequestration and aid the transition to a more sustainable natural resources management system and its monitoring. By integrating scenario-based modelled data with context-specific knowledge, this approach is likely to be a flexible solution to promote the transition, especially in a highly diverse sector such as agroforestry (Fischer and Glenk, 2011; Nicolosi and Feola, 2016).

Conclusion

Decisions to adopt or abandon silvopastoral systems to promote C sequestration are shaped by several factors (e.g. scale, profitability, labour requirements, trade-offs) shown to be significant in the agroecological transition process. Discussion of these factors centres on the role of scientific evidence and key categories, specifically, farming type, management practices and community characteristics. In addition, these factors are both challenges and opportunities for scientists, environmental professionals, farmers, land managers and policy makers to consider when planning and implementing agroecological systems such as agroforestry and silvopastoralism in particular. Hence it is suggested that the transition to agroecology could realistically be achieved by implementing a decision support tool where the benefits of agroforestry systems to store carbon can be incorporated and information and open communication links as main drivers are considered in adoption.

References

- Altieri, M.A. (2002) Agroecology: The science of natural resource management for poor farmers in marginal environments. *Agriculture, Ecosystems and Environment*, 93(1–3), 1–24. [http://doi.org/10.1016/S0167-8809\(02\)00085-3](http://doi.org/10.1016/S0167-8809(02)00085-3)
- Altieri, M.A., Nicholls, C.I., Henao, A. and Lana, M.A. (2015) Agroecology and the design of climate change-resilient farming systems. *Agronomy for Sustainable Development*. <http://doi.org/10.1007/s13593-015-0285-2>
- Biesbroek, G.R., Swart, R.J., Carter, T.R., Cowan, C., Henrichs, T., Mela, H., Morecroft, M.D. and Rey, D. (2010) Europe adapts to climate change: Comparing national adaptation strategies. *Global Environmental Change*, 20(3), 440-450.
- Barnes, A., Beechener, S., Cao, Y., Elliott, J., Harris, D., Jones, G., Toma, L. and Whiting, M. (2010) FF0201: Market segmentation in the agriculture sector: climate change. Final report to DEFRA.

- Bodin, Ö. and Tengö, M. (2012) Disentangling Intangible Social–Ecological Systems. *Global Environmental Change*, 22(2), 430–439.
- Black, K., Green, S., Mullooley, G. and Poveda, P. (2014) Carbon sequestration by hedgerows in the Irish landscape: Towards a National Hedgerow Biomass Inventory for the LULUCF Sector using LiDAR remote sensing. Climate Change Research Programme (CCRP) 2007-2013 Report Series No. 32. <http://www.epa.ie/pubs/reports/research/climate/ccrp-32-for-webFINAL.pdf>
- Bohan, D.A., Raybould, A., Mulder, C., Woodward, G., Tamaddoni-Nezhad, A., Bluthgen, N. ... Macfadyen, S. (2013) Networking agroecology. integrating the diversity of agroecosystem interactions. *Advances in Ecological Research* (Vol. 49). <http://doi.org/10.1016/B978-0-12-420002-9.00001-9>
- Broom, D.M., Galindo, F. and Murgueitio, E. (2013) Sustainable, efficient livestock production with high biodiversity and good welfare for animals. *Proceedings of the Royal Society B: Biological Sciences*, 280(1771), 20132025–20132025. <http://doi.org/10.1098/rspb.2013.2025>
- Burbi, S., Baines, R.N. and J.S. Conway (2016) Achieving successful farmer engagement on greenhouse gas emission mitigation. *International Journal of Agricultural Sustainability* 14: 466-483. <http://doi.org/10.1080/14735903.2016.1152062>
- Burbi, S. and Olave, R.J. (2017) Transition to agroforestry: current challenges and opportunities for the adoption of agroforestry as carbon sequestration strategy. Agroecology Forum 2017. 25-27 October 2017. ISARA, Lyon, France.
- Cardinael, R., Chevallier, T., Cambou, A., Béral, C., Barthès, B. G., Dupraz, C., ... and C. Chenu (2017) Increased soil organic carbon stocks under agroforestry: A survey of six different sites in France. *Agriculture, Ecosystems & Environment* 236, 243–255. <http://doi.org/10.1016/j.agee.2016.12.011>
- Castellanos, E.J., Tucker, C., Eakin, H., Morales, H., Barrera, J. F. and Díaz, R. (2013) Assessing the Adaptation Strategies of Farmers Facing Multiple Stressors: Lessons from the Coffee and Global Changes Project in Mesoamerica. *Environmental Science & Policy*, 26(0), 19–28.
- Cornell, S., Berkhout, F., Tuinstra, W., Tàbara, J. D., Jäger, J., Chabay, I., de Wit, B., Langlais, R., Mills, D., Moll, P., Otto, I.M., Petersen, A., Pohl, C. and van Kerkhoff, L. (2013) Opening Up Knowledge Systems for Better Responses to Global Environmental Change. *Environmental Science & Policy*, In Press, Corrected Proof, Available online 16 January 2013.
- Csikvari, J., Grandgirard, D., Reubens, B., Morhart, C.D., Pecenka, R., Schmutz, U., Vityi, A., Balaguer, F. and Considine W. (2017) MINIPAPER 3: Tools for Optimal Design and Management. EIP-Agri Focus Group. Agroforestry. 15 February 2017.
- de Boer, I., Cederberg, C., Eady, S., Gollnow, S., Kristensen, T., Macleod, M., Meul, M., Nemecek, T., Phong, L., Thoma, G., van der Werf, H., Williams, A. and Zonderland-Thomassen, M. (2011) Greenhouse gas mitigation in animal production: Towards an integrated life cycle sustainability assessment. *Current Opinion in Environmental Sustainability*, 3(5), 423-431.
- DEFRA (2016) 2016 Government GHG Conversion Factors for Company Reporting: Methodology Paper for Emission Factors. Department for Environment, Food and Rural Affairs and Department for Business, Energy & Industrial Strategy. 5 October 2016.
- den Herder, M., Moreno, G., Mosquera-Losada, R., Palma, J., Sidiropoulou, A., Javier, J., ... Burgess, P. (2017). Current extent and trends of agroforestry in the EU27. *Agriculture, Ecosystems and Environment* 241, 121–132. <http://doi.org/10.1016/j.agee.2017.03.005>
- Dumont B., González-García, E., Thomas, M., Fortun-Lamothe, L., Ducrot, C., Dourmad, J.Y. and Tichit, M. (2014) Forty research issues for the redesign of animal production systems in the 21st century *Animal* 8:8,1382–1393.
- Emery, S.B. and Franks, J.R. (2012) The Potential for Collaborative Agri-Environment Schemes in England: Can a Well-Designed Collaborative Approach Address Farmers' Concerns with Current Schemes?. *Journal of Rural Studies*, 28(3), 218–231.
- Fagerholm, N., Torralba, M., Burgess, P.J. and Plieninger, T. (2016) A systematic map of ecosystem services assessments around European agroforestry. *Ecological Indicators*, 62, 47–65. <http://doi.org/10.1016/j.ecolind.2015.11.016>
- Feola, G. and Binder, C.R. (2010) Towards an Improved Understanding of Farmers' Behaviour: The Integrative Agent-Centred (IAC) Framework. *Ecological Economics*, 69(12), 2323–2333.

- Fischer, A. and Glenk, K. (2011) One Model Fits all? — On the Moderating Role of Emotional Engagement and Confusion in the Elicitation of Preferences for Climate Change Adaptation Policies. *Ecological Economics*, 70(6), 1178–1188.
- Fornara, D., Olave, R., Burgess, P., Delmer, A., Upson, M. and J. McAdam (2017) Land use change and soil carbon pools: evidence from a long-term silvopastoral experiment. *Agroforestry Systems*, in Press. <http://doi.org/10.1007/s10457-017-0124-3>
- German, L.A., Tiani, A.M., Daoudi, A., Maravanyika, T.M., Chuma, E., Jum, C., Nemarundwe, N., Ontita, E. and Yitamben, G. (2012) The Application of Participatory Action Research to Climate Change Adaptation in Africa: a reference guide. Center for International Forestry Research (CIFOR), Bogor, Indonesia (2012) 106 pp.
- Hart, K. and Baldock, D. (2011) Greening the CAP: delivering environmental outcomes through pillar one. *Institute for European Environmental Policy*, 1(11), 1–26.
- Hofmann, M.E., Hinkel, J. and Wrobel, M. (2011) Classifying knowledge on climate change impacts, adaptation, and vulnerability in Europe for informing adaptation Research and decision-making: A conceptual meta-analysis. *Global Environmental Change*, 21(3), 1106-1116.
- Howlett, D.S., Moreno, G., Mosquera-Losada, M.R., Nair, P.K.R. and Nair, V.D. (2011) Soil carbon storage as influenced by tree cover in the Dehesa cork oak silvopasture of central-western Spain. *Journal of Environmental Monitoring: JEM*, 13(C), 1897–1904. <http://doi.org/10.1039/c1em10059a>
- IFAD (2009) Good practices in participatory mapping. A review prepared for the International Fund for Agricultural Development (IFAD). Editor: John Corbett, University of British Columbia, Okanagan: Canada. 59pp. <http://doi.org/9780132298346>
- Islam, M.M., Barnes, A. and Toma, L. (2013) An investigation into climate change scepticism among farmers. *Journal of Environmental Psychology*, 34(0), 137–150.
- Jacobi, J., Mathez-Stiefel, S.-L., Gambon, H., Rist, S. and Altieri, M. (2016) Use and Role of Local and External Knowledge in Agroforestry Projects in Bolivia. *Environmental Management*, 1–13. <http://doi.org/10.1007/s00267-016-0805-0>
- Jones, A.K., Jones, D.L., Edwards-Jones, G and Cross, P. (2013) Informing decision-making in agricultural greenhouse gas mitigation policy: A Best–Worst Scaling survey of expert and farmer opinion in the sheep industry. *Environmental Science & Policy*. Article In Press.
- Keesman, K.J., Graves, A., van der Werf, W., Burgess, P.J., Palma, J., Dupraz, C. and van Keulen, H. (2011) A system identification approach for developing and parameterising an agroforestry system model under constrained availability of data. *Environmental Modelling and Software*, 26(12), 1540–1553. <http://doi.org/10.1016/j.envsoft.2011.07.020>
- Khatri-Chhetri, A., Aggarwal, P.K., Joshi, P.K. and Vyas, S. (2017) Farmers' prioritization of climate-smart agriculture (CSA) technologies. *Agricultural Systems*, 151, 184–191. <http://doi.org/10.1016/j.agsy.2016.10.005>
- Kings, D. and Ilbery, B. (2010) The environmental belief systems of organic and conventional farmers: Evidence from central-southern England. *Journal of Rural Studies*, 26(4), 437–448.
- Klerkx, L., Schut, M., Leeuwis, C. and Kilelu, C. (2012) Advances in Knowledge Brokering in the Agricultural Sector: Towards Innovation System Facilitation. Institute of Development Studies, Oxford, England. IDS Bulletin. Volume 43, Number 5. September 2012.
- Kumar, B.M. and Nair, P.K.R. (2011) Carbon sequestration in European agroforestry systems - Opportunities and Challenges. *Advances in Agroforestry*. Springer. <http://doi.org/10.1007/978-94-007-1630-8>
- Le Gal, P.-Y., Dugué, P., Faure, G. and Novak, S. (2011) How does Research Address the Design of Innovative Agricultural Production Systems at the Farm Level? A Review. *Agricultural Systems*, 104(9), 714–728.
- Llewellyn, R.S. (2007) Information quality and effectiveness for more rapid adoption decisions by farmers. *Field Crops Research*, 104(1–3), 148-156.
- Long, T.B., Blok, V. and Coninx, I. (2016) Barriers to the adoption and diffusion of technological innovations for climate-smart agriculture in Europe: Evidence from the Netherlands, France, Switzerland and Italy. *Journal of Cleaner Production*, 112, 9–21. <http://doi.org/10.1016/j.jclepro.2015.06.044>

- Louah, L., Visser, M., Blaimont, A. and de Cannière, C. (2017) Barriers to the development of temperate agroforestry as an example of agroecological innovation: Mainly a matter of cognitive lock-in? *Land Use Policy*, 67, 86–97. <http://doi.org/10.1016/j.landusepol.2017.05.001>
- Lovell, S.T., DeSantis, S., Nathan, C.A., Breton Olson, M., Méndez, V.E., Kominami, H. C., Erikson, D.L., Morris, K.S. and Morris, W.B. (2010) Integrating agroecology and landscape multifunctionality in Vermont: An evolving framework to evaluate the design of agroecosystems. *Agricultural Systems*, 103(5), 327–341. <http://doi.org/10.1016/j.agsy.2010.03.003>
- Luedeling, E., Kindt, R., Huth, N.I. and Koenig, K. (2014) Agroforestry systems in a changing climate- Challenges in projecting future performance. *Current Opinion in Environmental Sustainability*, 6(1), 1–7. <http://doi.org/10.1016/j.cosust.2013.07.013>
- Mapfumo, P., Adjei-Nsiah, S., Mtambanengwe, F., Chikowo, R. and Giller, K.E. (2013) Participatory Action Research (PAR) as an Entry Point for Supporting Climate Change Adaptation by Smallholder Farmers in Africa. *Environmental Development*, 5(0), 6–22.
- McKenzie F. (2011) Farmer-driven Innovation in New South Wales, Australia. *Australian Geographer*, 44(1), 81–95.
- Mills, J., Gaskell, P., Reed, M., Short, C., Ingram, J., Boatman, N., Jones, N., Conyers, S., Carey, P., Winter, M. and Lobley, M. (2013) Farmer attitudes and evaluation of outcomes to on-farm environmental management. Report to Department for Environment, Food and Rural Affairs (Defra). CCRI: Gloucester.
- Mosquera-Losada, M.R., Borek, R., Balaguer, F., Mezzarila, G. and Ramos-Font, M.E. (2017) MINIPAPER 9: Agroforestry as a mitigation and adaptation tool. EIP-Agri Focus Group. Agroforestry. 25 February 2017. https://ec.europa.eu/eip/agriculture/sites/agri-eip/files/fg22_mp9_cc_adaptation_mitigation_2017_en.pdf
- Mosquera-Losada, M.R., Moreno, G., Santiago-Freijanes, J.J., Ferreiro-Domínguez, N. and Rigueiro-Rodríguez, A. (2015) Sistemas agroforestales y PAC. *Ambienta*, 112(SEPTEMBER), 110–124.
- Mwongera, C., Shikuku, K.M., Twyman, J., Läderach, P., Ampaire, E., Van Asten, P., Twomlow, S. and Winowiecki, L.A. (2017) Climate smart agriculture rapid appraisal (CSA-RA): A tool for prioritizing context-specific climate smart agriculture technologies. *Agricultural Systems*, 151, 192–203. <http://doi.org/10.1016/j.agsy.2016.05.009>
- Neef, A. and Neubert, D. (2011) Stakeholder Participation in Agricultural Research Projects: A Conceptual Framework for Reflection and Decision-making'. *Agriculture and Human Values* 28,179–94.
- Ness, B., Urbel-Piirsalu, E., Anderberg, S. and Olsson, L. (2007) Categorising tools for sustainability assessment. *Ecological Economics*, 60(3), 498–508. doi:[10.1016/j.ecolecon.2006.07.023](https://doi.org/10.1016/j.ecolecon.2006.07.023)
- Newell, B., Crumley, C.L., Hassan, N., Lambin, E.F., Pahl-Wostl, C., Underdal, A. and Wasson, R. (2005) A Conceptual Template for Integrative human–environment Research. *Global Environmental Change*, 15(4), 299–307.
- Nguyen F., Doreau, M., Eugene, M., Corson, M., Garcia-Launay, F., Chesneau, G and van der Werf, H. (2013) Effect of farming practices for greenhouse gas mitigation and subsequent alternative land use on environmental impacts of beef cattle production systems. *Animal*, 7, 860-869.
- Nicholls, C.I, Altieri, M.A., Dezanet, A., Lana, M., Feistauer, D. and Ouriques, M. (2014) A rapid, farmer-friendly agroecological method to estimate soil quality and crop health in vineyard systems. *Sci. Ecol. Biodynamics*, Autumn, 33–40.
- Nicolosi, E. and Feola, G. (2016) Transition in place: Dynamics, possibilities, and constraints. *Geoforum*, 76(November), 153–163. <http://doi.org/10.1016/j.geoforum.2016.09.017>
- Olave, R. (2016) Agroforestry as a land use option to sequester carbon in a cool temperate climate. World Congress Silvo-pastoral Systems 2016. University of Évora, Portugal. 27-30 September 2016.
- Oliver, D.M., Fish, R.D., Winter, M., Hodgson, C.J., Heathwaite, A.L. and Chadwick, D.R. (2012) Valuing Local Knowledge as a Source of Expert Data: Farmer Engagement and the Design of Decision Support Systems. *Environmental Modelling & Software*, 36(0), 76–85.
- Oreszczyn, S., Lane, A. and Carr, S. (2010) The Role of Networks of Practice and Webs of Influencers on Farmers' Engagement with and Learning about Agricultural Innovations. *Journal of Rural Studies*, 26(4), 404–417.

- Palma, J.H.N., Crous-Duran, J., Graves, A.R. and Burgess, P.J. (2015) Database of Agroforestry System Descriptions. Milestone Report 28 (6.3) for EU FP7 Research Project: AGFORWARD 613520. (20 October 2015). 81 pp.
- Paracchini, M.L., Petersen, J.-E., Hoogeveen, Y., Bamps, C., Burfield, I. and Swaay, C. van (2008) *High Nature Value Farmland in Europe - An Estimate of the Distribution Patterns on the Basis of Land Cover and Biodiversity Data*. Institute for Environment and Sustainability Office for Official Publications of the European Communities Luxembourg. <http://doi.org/10.2788/8891>
- Pretty, J. and Buck, L.E. (2002) Social Capital and Social Learning in the Process of Natural Resource Management. In: C.B. Barrett, F. Place and A.A. Aboud (eds.), *Natural Resources Management in African Agriculture*. Nairobi: ICRAF and CABI Publishing.
- Pretty, J., Toulmin, C. and Williams, S. (2011) Sustainable intensification in African agriculture. *International Journal of Agricultural Sustainability* 9(1), 5-24.
- Raymond, C.M., Fazey, I., Reed, M.S., Stringer, L.C., Robinson, G.M. and Evely, A.C. (2010) Integrating Local and Scientific Knowledge for Environmental Management. *Journal of Environmental Management*, 91(8), 1766–1777.
- Rakotovo, N.H., Razafimbelo, T.M., Rakotosamimanana, S., Randrianasolo, Z., Randriamalala, J.R. and Albrecht, A. (2017) Carbon footprint of smallholder farms in Central Madagascar: The integration of agroecological practices. *Journal of Cleaner Production*, 140, 1165–1175. <http://doi.org/10.1016/j.jclepro.2016.10.045>
- Rittenhouse, C.D. and Rissman, A.R. (2012) Forest cover, carbon sequestration, and wildlife habitat: Policy review and modeling of tradeoffs among land-use change scenarios. *Environmental Science and Policy*, 21, 94–105. <http://doi.org/10.1016/j.envsci.2012.04.006>
- Saj, S., Torquebiau, E., Hainzelin, E., Pages, J. and Maraux, F. (2017) The way forward: An agroecological perspective for Climate-Smart Agriculture. *Agriculture, Ecosystems and Environment*, 250, 20–24. <http://doi.org/10.1016/j.agee.2017.09.003>
- Sarandón, S. and Flores, C. (2014) *Agroecología: bases teóricas para el diseño y manejo de agroecosistemas sustentables*. <http://doi.org/E-Book>
- Schöll, R. and Binder, C. (2009) Comparing system visions of farmers and experts. *Futures*, 41(9), 631-649.
- Schut, M., Klerkx, L., Sartas, M., Lamers, D., Campbell, M.M.C., Ogbonna, I., Kaushik, P., Atta-Krah, K. and Leeuwis, C. (2015) Innovation Platforms: Experiences with Their Institutional Embedding in Agricultural Research for Development. *Experimental Agriculture*, 1–25. <http://doi.org/10.1017/S001447971500023X>
- Sereke, F., Graves, A.R. and Herzog, F. (2016) Drivers of Swiss agroforestry it's not all about money. 3rd European Agroforestry conference 2016, Montpellier, France, 23 -25 May 2016.
- Sutherland, L.A., Burton, R.J.F., Ingram, J., Blackstock, K., Slee, B. and Gotts, N. (2012) Triggering change: Towards a conceptualisation of major change processes in farm decision-making. *Journal of Environmental Management*, 104, 142–151. <http://doi.org/10.1016/j.jenvman.2012.03.013>
- Swanson, B.E. and Rajalahti, R. (2010) Strengthening agricultural extension and advisory systems: Procedures for assessing, transforming and evaluating extension systems. World Bank Agriculture and Rural Development discussion paper 45. Washington DC: IBRD/World Bank.
- Takemura, K., Uchida, Y. and Yoshikawa, S. (2014) Roles of Extension Officers to Promote Social Capital in Japanese Agricultural Communities. *PLoS ONE* 9(3): e91975.
- Tittonell, P., Scopel, E., Andrieu, N., Posthumus, H., Mapfumo, P., Corbeels, M., van Halsema, G.E., Lahmar, R., Lugandu, S., Rakotoarisoa, J., Mtamanengwe, F., Pound, B. Chikowo, R., Naudin, K., Triomphe, B. and Mkomwa, S. (2012) Agroecology-based aggradation-conservation agriculture (ABACO): Targeting innovations to combat soil degradation and food insecurity in semi-arid Africa. *Field Crops Research*, 132, 168–174. <http://doi.org/10.1016/j.fcr.2011.12.011>
- Torralba, M., Fagerholm, N., Burgess, P.J., Moreno, G. and Plieninger, T. (2016) Do European agroforestry systems enhance biodiversity and ecosystem services? A meta-analysis. *Agriculture, Ecosystems and Environment*, 230, 150–161. <http://doi.org/10.1016/j.agee.2016.06.002>

- TWN and SOCLA (2015) *Agroecology: Key Concepts, Principles and Practices*. Third World Network (TWN) and Sociedad Científica Latinoamericana de Agroecología. SOCLA. 54pp.
- UNFCCC (2015) Adoption of the Paris agreement – proposal by the President – draft decision. FCCC/CP/ 2015/L.9. 21st Conference of the Parties of the United Nations Framework Convention on Climate Change (UNFCCC). 30 November – 12 December 2015 Paris, France.
- United Kingdom Parliament (2008) UK Climate Change Act 2008. Chapter 27. London, United Kingdom.
- Vellinga, T.V., de Haan, M.H.A., Schils, R.L.M., Evers, A. and van den Pol–van Dasselaar, A. (2011) Implementation of GHG mitigation on intensive dairy farms: Farmers' preferences and variation in cost effectiveness. *Livestock Science*, 137(1–3), 185-195.
- Virji, H., Padgham, J. and Seipt, C. (2012) Capacity building to support knowledge systems for resilient development — approaches, actions, and needs. *Current Opinion in Environmental Sustainability*, 4(1), 115–121.
- Wallach, D., Mearns, L.O., Ruane, A.C., Rötter, R.P. and Asseng, S. (2016) Lessons from climate modelling on the design and use of ensembles for crop modelling. *Climatic Change*, 139(3–4), 551–564. <http://doi.org/10.1007/s10584-016-1803-1>
- Wezel, A., Bellon, S., Doré, T., Francis, C., Vallod, D. and David, C. (2009) Agroecology as a science, a movement and a practice. A review. *Agronomy for Sustainable Development*, 1–13. <http://doi.org/10.1051/agro/2009004>
- Wezel, A., Casagrande, M., Celette, F., Vian, J.F., Ferrer, A. and Peigné, J. (2014) Agroecological practices for sustainable agriculture. A review. *Agronomy for Sustainable Development*. <http://doi.org/10.1007/s13593-013-0180-7>
- Wibbelmann, M., Schmutz, U., Wright, J., Udall, D., Rayns, F., Kneafsey, M., Trenchard, L., Bennett, J. and Lennartsson, M. (2013) *Mainstreaming Agroecology: Implications for Global Food and Farming Systems*. Centre for Agroecology and Food Security Discussion Paper. Coventry: Centre for Agroecology and Food Security. ISBN: 978-1-84600-0454
- Wilson, M.H. and S.T. Lovell (2016) Agroforestry-The next step in sustainable and resilient agriculture. *Sustainability* 8(6), 1–15. <http://doi.org/10.3390/su8060574>
- Xu, J.C., Kim, G.J. and He, J. (2011) Participatory Agroforestry Development in DPR Korea. Kunming: World Agroforestry Centre (ICRAF), China and East Asia Node.